

On rack polynomials

Tim Carrell

Sam Nelson

Abstract

We study rack polynomials and the link invariants they define. We show that constant action racks are classified by their generalized rack polynomials and show that ns^at^a -quandles are not classified by their generalized quandle polynomials. We use subrack polynomials to define enhanced rack counting invariants, generalizing the quandle polynomial invariants.

KEYWORDS: Finite racks, rack polynomials, knot and link invariants
2000 MSC: 57M25, 57M27, 17D99

1 Introduction

In [6], a two-variable polynomial invariant of finite quandles was introduced. This polynomial quantifies the way in which the trivial action of one quandle element on another is distributed throughout the quandle as opposed to concentrated in a single identity element as in a group.

In [7] the quandle polynomial was generalized to a family of N^2 polynomials where N is the least common multiple of the exponents of the columns of the quandle matrix considered as elements of the symmetric group S_n on the elements of the quandle. In both cases, the quandle polynomials were used to enhance the quandle counting invariants to obtain new invariants which specialize to the original quandle counting invariants but contain more information.

In this paper we study the natural generalization of quandle polynomials to finite racks. We are able to show that for at least one class of finite racks, the generalized rack polynomials determine the rack structure up to isomorphism, and we identify another class of quandles for which the generalized rack polynomials do not determine the isomorphism class. We then use these rack polynomials to enhance the rack counting invariants from [8].

The paper is organized as follows. In section 2 we review the definitions of racks and rack polynomials and give some examples. In section 3 we show that constant action racks are classified by their generalized rack polynomials. In section 4 we show that Alexander quandles have quandle polynomial of the form ns^at^a and that unlike constant action racks, Alexander quandles and other quandles with quandle polynomial ns^at^a are not classified by their generalized quandle polynomials. In section 5 we define rack polynomial enhanced counting invariants. In section 6 we collect some questions for future investigation.

2 Racks and rack polynomials

In [4], Joyce defined a kind of self-distributive algebraic structure which he dubbed a “quandle.” In [3], quandles were generalized to a larger class of self-distributive algebraic systems known as “racks.” Both concepts appear under other names in the literature such as “distributive groupoids,” “automorphic sets” and “kei.” See [5, 2, 11].

Definition 1 A *rack* is a set X with a binary operation $\triangleright : X \times X \rightarrow X$ satisfying

- (i) for all $x, y \in X$ there is a unique $z \in X$ satisfying $x = z \triangleright y$, and
- (ii) for all $x, y, z \in X$ we have $(x \triangleright y) \triangleright z = (x \triangleright z) \triangleright (y \triangleright z)$.

A rack which additionally satisfies

$$(0) \text{ for all } x \in X, \text{ we have } x \triangleright x = x$$

is a *quandle*.

Axiom (i) requires that each element x of a rack X acts on X bijectively, while axiom (ii) requires these bijections to be automorphisms of the rack structure. The bijectivity of the action of x gives us a right inverse action $\triangleright^{-1} : X \times X \rightarrow X$ defined by $x \triangleright^{-1} y = z$ where $x = z \triangleright y$. The reader can check that $(X, \triangleright^{-1})$ is also a rack, called the *dual* of $(X, \triangleright)$.

Standard examples of rack and quandle structures include:

- any union of conjugacy classes in a group G with operation $a \triangleright b = b^{-n}ab^n$, $n \in \mathbb{Z}$
- the set of right cosets in a group G of a subgroup $H \subset G$ (not necessarily normal) fixed by an automorphism $s : G \rightarrow G$ with rack operation $Hx \triangleright Hy = s(HxHy^{-1})Hy$
- any set X with a permutation $\sigma \in S_X$ with $x \triangleright y = \sigma(x)$ (these are *constant action racks* or *permutation racks*)
- the subset of a vector space V on which a bilinear form $\langle \mathbf{x}, \mathbf{x} \rangle \neq 0$ with

$$\mathbf{x} \triangleright \mathbf{y} = \alpha \left(\mathbf{x} - 2 \frac{\langle \mathbf{x}, \mathbf{y} \rangle}{\langle \mathbf{y}, \mathbf{y} \rangle} \mathbf{y} \right)$$

where α is a non-zero scalar (these are called *Coxeter racks*; see [3, 9])

- any module over $\mathbb{Z}[t^{\pm 1}, s]/s(1-t-s)$ with $x \triangleright y = tx + sy$.

Racks of the last type in which $s = 1 - t$ are known as *Alexander quandles*. A rack is *abelian* if for all $x, y, z, w \in X$ we have

$$(x \triangleright y) \triangleright (z \triangleright w) = (x \triangleright z) \triangleright (y \triangleright w).$$

In addition to being right-distributive, abelian quandles are also left-distributive, since we have

$$a \triangleright (b \triangleright c) = (a \triangleright a) \triangleright (b \triangleright c) = (a \triangleright b) \triangleright (a \triangleright c).$$

A quandle is a *crossed set* (see [1]) if we have

$$x \triangleright y = x \iff y \triangleright x = y.$$

The reader can check that Alexander quandles are abelian and Coxeter quandles (set $\alpha = -1$) are crossed sets.

We can express rack structures on a finite set $X = \{x_1, \dots, x_n\}$ in an algebra-agnostic way, i.e. without needing a formula for $x \triangleright y$, by giving the rack operation table as a matrix M_X whose (i, j) entry is k where $x_k = x_i \triangleright x_j$. We call this the *rack matrix* of X .

Example 1 The constant action rack on $X = \{1, 2, 3\}$ with $\sigma = (132)$ has rack matrix

$$M_X = \begin{bmatrix} 3 & 3 & 3 \\ 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix}.$$

For a rack X , say that an equivalence relation $\sim$ on X is a *congruence* if $x \sim x'$ and $y \sim y'$ imply $x \triangleright y \sim x' \triangleright y'$. The set $X/\sim$ of equivalence classes then forms a *quotient rack* under the operation $[x \triangleright y] = [x] \triangleright [y]$. See [10] for more.

Next, we have a definition from [7]:

Definition 2 Let X be a finite rack. For each $x \in X$, define

$$C_m(x) = \{y \in X \mid y \triangleright^m x = y\} \quad \text{and} \quad R_n(x) = \{y \in X \mid x \triangleright^n y = x\}$$

where

$$x \triangleright^i y = (\dots (x \triangleright y) \triangleright y) \dots \triangleright y$$

where i is the number of triangles. Denote $c_m(x) = |C_m(x)|$ and $r_n(x) = |R_n(x)|$. Then the (m, n) -rack polynomial of X (or (m, n) -quandle polynomial if X is a quandle) is

$$rp_{m,n}(X) = \sum_{x \in X} s^{c_m(x)} t^{r_n(x)}.$$

The terms “rack polynomial” and “quandle polynomial” without specified m and n values will refer to the case $m = n = 1$.

Example 2 The constant action rack X with rack matrix $M_X = \begin{bmatrix} 2 & 2 & 2 \\ 1 & 1 & 1 \\ 3 & 3 & 3 \end{bmatrix}$ has rack polynomial $rp_{1,1}(X) = 2t + s^3t$.

Remark 3 In [6] example 8, it is incorrectly stated that a rack may have rack polynomial equal to zero, since in that example we have a contribution of s^0t^0 from each element. Of course, $s^0t^0 = 1 \neq 0$, and indeed the coefficients of a rack polynomial always sum to the cardinality $|X|$. The second listed author is grateful to the first for catching this oversight.

3 Generalized rack polynomials of constant action racks

In this section we show that constant action racks are classified by their generalized rack polynomials.

Proposition 1 Let X be the constant action rack of a given permutation σ on $\{x_1, x_2, \dots, x_k\}$. Then the generalized rack polynomial of X is

$$rp_{m,n}(X) = bs^k t^a + (k - b)t^a$$

where a is the number of x_i such that $\sigma^n(x_i) = x_i$ and b is the number of x_i such that $\sigma^m(x_i) = x_i$.

Proof. For any $x \in \{x_1, x_2, \dots, x_k\}$,

$$c_n(x) = |\{y \mid y \triangleright^n x = y\}| = |\{y \mid \sigma^n(y) = y\}| = a$$

so each term of $rp_{m,n}(X)$ contains a factor of t^a . Furthermore,

$$r_m(x) = |\{y \mid x \triangleright^m y = x\}| = |\{y \mid \sigma^m(x) = x\}| = \begin{cases} 0 & \text{if } \sigma^m(x) \neq x, \\ k & \text{if } \sigma^m(x) = x. \end{cases}$$

Therefore, each x such that $\sigma^m(x) = x$ corresponds to a term of $s^k t^a$, and each x such that $\sigma^m(x) \neq x$ corresponds to a term of t^a . There are b distinct x such that $\sigma^m(x) = x$, and thus $k - b$ distinct x such that $\sigma^m(x) \neq x$. Therefore,

$$rp_{m,n}(X) = bs^k t^a + (k - b)t^a.$$

□

Note that the original “non-generalized” rack polynomial corresponds to $rp_{1,1}(X)$. For this specific case, we have a and b both equal to the number of fixed points of σ , and so we have

Corollary 2 *Let X be the constant action rack of a given permutation σ on $\{x_1, x_2, \dots, x_k\}$, and suppose σ has b fixed points. Then the rack polynomial of X is*

$$rp(X) = bs^k t^b + (k - b)t^b.$$

Proposition 3 *The set of generalized rack polynomials is a complete invariant of constant action racks.*

To prove this result we will need a pair of lemmas.

Lemma 4 *Suppose X and X' are constant action racks given by permutations σ and σ' , respectively. Then X and X' are isomorphic if and only if σ and σ' have the same cycle structure.*

Proof. Suppose that X and X' are isomorphic. Then for any positive integer l ,

$$rp_{l,1}(X) = rp_{l,1}(X')$$

and thus X and X' must have the same number of cycles whose length divides l . For $l = 1$, this means that X and X' have the same number of cycles of length 1. Proceeding inductively, X and X' have the same number of cycles of all positive integer lengths l , and so have the same cycle structure.

Conversely, suppose that σ and σ' have the same cycle structure, that is, we can write them as

$$\sigma = (1_1 1_2 \dots 1_{n_1})(2_1 2_2 \dots 2_{n_2}) \dots (k_1 k_2 \dots k_{n_k})$$

and

$$\sigma' = (1'_1 1'_2 \dots 1'_{n_1})(2'_1 2'_2 \dots 2'_{n_2}) \dots (k'_1 k'_2 \dots k'_{n_k}).$$

Let $\alpha : \sigma \rightarrow \sigma'$ be the bijection given by $a_i \mapsto a'_i$. Then

$$\alpha(a_i \triangleright b_j) = \alpha(\sigma(a_i)) = \alpha(a_{i+1}) = a'_{i+1} = \sigma'(a'_i) = a'_i \triangleright b'_j = \alpha(a_i) \triangleright \alpha(b_j)$$

and so α is a rack isomorphism from X to X' . □

Lemma 5 *Suppose X and X' are constant action racks given by the permutations σ and σ' , respectively. Then $rp_{m,n}(X) = rp_{m,n}(X')$ for all $m, n \in \mathbb{Z}^+$ if and only if σ and σ' have the same cycle structure.*

Proof. By proposition 1, if X and X' are constant action racks of the same cardinality, then $rp_{m,n}(X) = rp_{m,n}(X')$ if and only if

$$|\{x_i \mid \sigma^m(x_i) = x_i\}| = |\{x_i \mid \sigma'^m(x_i) = x_i\}| \quad (1)$$

and

$$|\{x_i \mid \sigma^n(x_i) = x_i\}| = |\{x_i \mid \sigma'^n(x_i) = x_i\}|. \quad (2)$$

If σ and σ' have the same cycle structure, then they have the same number of elements in cycles whose length divides m and the same number of elements in cycles whose length divides n . Thus, both (1) and (2) hold and so $rp_{m,n}(X) = rp_{m,n}(X')$.

If σ and σ' do not have the same cycle structure, then there exists some minimal l such that σ and σ' do not have the same number of cycles of length l . Thus, for $m = l$ (1) cannot hold, and so $rp_{m,n}(X) \neq rp_{m,n}(X')$. □

Combining lemmas 4 and 5 immediately gives us proposition 3. The following example shows that the generalized rack polynomials must be used to get a complete invariant on constant action racks; the $(1, 1)$ -rack polynomial is not sufficient.

Example 4 The racks with rack matrices

$$M_X = \begin{bmatrix} 2 & 2 & 2 & 2 & 2 & 2 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 4 & 4 & 4 & 4 & 4 & 4 \\ 3 & 3 & 3 & 3 & 3 & 3 \\ 6 & 6 & 6 & 6 & 6 & 6 \\ 5 & 5 & 5 & 5 & 5 & 5 \end{bmatrix} \quad \text{and} \quad M_Y = \begin{bmatrix} 2 & 2 & 2 & 2 & 2 & 2 \\ 3 & 3 & 3 & 3 & 3 & 3 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 5 & 5 & 5 & 5 & 5 & 5 \\ 6 & 6 & 6 & 6 & 6 & 6 \\ 4 & 4 & 4 & 4 & 4 & 4 \end{bmatrix}$$

both have no fixed points and so by corollary 2 have the same rack polynomial $rp(X) = rp(Y) = 6$, but have different cycle structures and so by lemma 4 are not isomorphic.

4 Quandle Polynomials of Alexander Quandles

In this section we study the quandle polynomials of Alexander quandles.

Definition 3 Let Q be an Alexander quandle. Say that x and y are $(1-t)$ -equivalent, denoted $x \sim_{(1-t)} y$ or just $x \sim y$, if $(1-t)x = (1-t)y$.

Proposition 6 Let Q be an Alexander quandle such that $|Q| = n$. Then

$$rp(Q) = ns^a t^a$$

for some positive integer $a|n$.

Proof. Let $\varphi : Q \rightarrow Q$ be the function given by

$$\varphi(q) = 0 \triangleright q.$$

Since

$$0 \triangleright q = t \cdot 0 + (1-t) \cdot q = (1-t) \cdot q,$$

φ is just left multiplication by $1-t$. Since Alexander quandles are abelian and thus left distributive, φ is automatically a homomorphism. By the first isomorphism theorem φ partitions Q into cosets by the congruence

$$q \sim p \iff \varphi(q) = \varphi(p).$$

These cosets are all of size $a = |\ker \varphi|$; clearly $a|n$. We will show that $q \triangleright p = q$ if and only if $q \sim p$.

Suppose $q \triangleright p = q$. Then

$$\begin{aligned} tq + (1-t)p &= q \\ (t-1)q + (1-t)p &= 0 \\ (1-t)(p-q) &= 0 \end{aligned}$$

so $p-q \in \ker \varphi$ and so $q \sim p$.

Suppose $q \sim p$. Then $0 = \varphi(p-q)$ implies

$$0 = \varphi_0(p-q) = (1-t)(p-q) = (t-1)q + (1-t)p$$

and hence

$$q = tq + (1-t)p = q \triangleright p.$$

Therefore, for any $q \in Q$ there are exactly a choices of p such that $q \triangleright p = q$ and also exactly a choices of p such that $p \triangleright q = p$. Thus, $r(q) = c(q) = a$ for all $q \in Q$, and so

$$rp(Q) = ns^a t^a.$$

□

Note that Proposition 6 provides a second, equivalent definition for $(1-t)$ -equivalence.

Corollary 7 *An equivalent definition of $\sim$ is $p \sim q$ if and only if $p \triangleright q = p$.*

This also gives us another proof of the fact noted in [1] that

Corollary 8 *All Alexander quandles are crossed sets.*

Proof. This follows immediately from corollary 7 and the reflexivity of equivalence relations. $\square$

These results suggest the possibility of defining $\sim$ for all crossed sets, or proving the converse of 6. However, there is a counterexample to both of these natural conjectures.

Example 5 Let Q be the crossed set with matrix

$$M_Q = \begin{bmatrix} 1 & 3 & 2 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 & 2 & 2 \\ 2 & 1 & 3 & 3 & 3 & 3 \\ 4 & 4 & 4 & 4 & 6 & 5 \\ 5 & 5 & 5 & 6 & 5 & 4 \\ 6 & 6 & 6 & 5 & 4 & 6 \end{bmatrix}.$$

Suppose, as for Alexander quandles and ns^at^a quandles, we define $\sim$ by $p \sim q$ when $p \triangleright q = p$. Then for Q we have $1 \sim 4$ and $4 \sim 2$, but $1 \not\sim 2$. Therefore, $\sim$ cannot be an equivalence relation for this crossed set.

Furthermore,

$$(1 \triangleright 1) \triangleright (4 \triangleright 2) = 1 \triangleright 4 = 1,$$

but

$$(1 \triangleright 4) \triangleright (1 \triangleright 2) = 1 \triangleright 3 = 2$$

and so Q is not abelian and thus not Alexander, disproving the converse of proposition 6.

This also provides a counterexample to the conjecture in [7] that distinct non-Latin quandles are distinguished by at least one of their generalized rack polynomials.

Example 6 Let R be the quandle with quandle matrix

$$M_R = \begin{bmatrix} 1 & 1 & 2 & 2 & 1 & 1 \\ 2 & 2 & 1 & 1 & 2 & 2 \\ 3 & 3 & 3 & 3 & 4 & 4 \\ 4 & 4 & 4 & 4 & 3 & 3 \\ 6 & 6 & 5 & 5 & 5 & 5 \\ 5 & 5 & 6 & 6 & 6 & 6 \end{bmatrix}.$$

Both R and Q from example 5 have the generalized rack polynomial

$$rp_{(m,n)}(R) = rp_{(m,n)}(Q) = 6s^c t^d$$

with

$$c = \begin{cases} 6 & \text{when } n \text{ is even,} \\ 4 & \text{when } n \text{ is odd,} \end{cases} \quad \text{and} \quad d = \begin{cases} 6 & \text{when } m \text{ is even,} \\ 4 & \text{when } m \text{ is odd.} \end{cases}$$

However, R is abelian and Q is not abelian, so Q and R are not isomorphic.

5 Rack polynomial enhanced link invariants

In [8], the quandle counting invariant $|\text{Hom}(Q(L), T)|$ was extended to the case of finite non-quandle racks. In this section we will enhance this invariant with rack polynomials. We begin by recalling how this was done in the quandle case.

Definition 4 Let S be a subrack $S \in X$. The (m, n) -subrack polynomial is

$$srp_{S \subset X}^{m,n}(s, t) = \sum_{x \in S} s^{c_m(x)} t^{r_n(x)}.$$

The subquandle polynomials of the image subquandles in $\text{Hom}(Q(L), T)$ are used to enhance the quandle counting invariants in [6, 7]. Specifically, instead of counting 1 for each element of $\text{Hom}(Q(L), T)$ to obtain the quandle counting invariant $|\text{Hom}(Q(L), T)|$, we count $srp_{\text{Im}(f) \subset T}^{m,n}(s, t)$ to obtain a multiset of subquandle polynomials. We can express these multisets in a polynomial-style form by writing the elements of the multiset as powers of a variable z and the multiplicities as coefficients.

Definition 5 Let L be a link and T a finite quandle. The (m, n) -subquandle polynomial invariant of L with respect to T is then

$$sp_{m,n}(L, T) = \sum_{f \in \text{Hom}(Q(L), T)} z^{srp_{\text{Im}(f) \subset T}^{m,n}(s, t)}.$$

Now, let L be an oriented link with c ordered components. For any diagram D of L , we can regard D as a framed link using the blackboard framing, i.e. giving each component of L a framing number w_i equal to its self-writhe. Thus, such a diagram has a framing vector $\mathbf{w} = (w_1, \dots, w_c) \in \mathbb{Z}^c$.

For any finite rack T , let $N(T)$ be the rack rank of T , i.e. the exponent of the permutation in $S_{|T|}$ along the diagonal of the rack matrix of T . If two ambient isotopic diagrams of D have writhe vectors which are componentwise congruent modulo $N(T)$, then there is a bijection

$$\phi : \text{Hom}(FR(D, \mathbf{w}), T) \rightarrow \text{Hom}(FR(D, \mathbf{w}'), T)$$

between the sets of rack homomorphisms from the fundamental racks of $(D, \mathbf{w})$ and $(D, \mathbf{w}')$ into T defined by sending a coloring of one diagram to a coloring of the same diagram with mN kinks added. Indeed, since any subrack containing an element $x \in T$ must also contain the rack powers $x^{\triangleright n}$ for all $n \in \mathbb{Z}$ (see [8]), ϕ preserves image subracks. Hence, as far as T is concerned, the framing vectors of D live in $W = (\mathbb{Z}_{N(T)})^c$, and we have an invariant of unframed links given by

$$SR(L, T) = |\{f \in \text{Hom}(FR(D, \mathbf{w}), T) \mid \mathbf{w} \in W\}|,$$

called the *simple rack counting invariant*. A refinement obtained by keeping track of which colorings belong to which framings is the *rack counting polynomial* invariant

$$PR(L, T) = \sum_{\mathbf{w} \in W} \left(|\text{Hom}(FR(D, \mathbf{w}), T)| \prod_{i=1}^c q_i^{w_i} \right).$$

When T is a quandle, $N(T) = 1$ and we have $SR(L, T) = |PR(L, T)| = |\text{Hom}(Q(L), T)|$. See [8] for more.

We would like to jazz up these rack counting invariants with the generalized rack polynomials. To this end, we propose the following

Definition 6 Let L be a link of c components, T a finite rack with rack rank $N(T)$, and $W = (\mathbb{Z}_{N(T)})^c$. Then the (m, n) -simple subrack polynomial enhanced rack counting multiset is the multiset

$$srpm_{m,n}(L, T) = \left\{ srp_{\text{Im}(f) \subset T}^{m,n}(s, t) \mid \mathbf{w} \in W, f \in \text{Hom}(FR(D, \mathbf{w}), T) \right\}$$

and the (m, n) -subrack polynomial enhanced rack counting multiset is the multiset of ordered pairs

$$rpm_{m,n}(L, T) = \left\{ \left(srp_{\text{Im}(f) \subset T}^{m,n}(s, t), \prod_{i=1}^c q_i^{w_i} \right) \mid \mathbf{w} \in W, f \in \text{Hom}(FR(D, \mathbf{w}), T) \right\}$$

We can also define the invariants in a more polynomial-style form for ease of comparison:

$$srpp_{m,n}(L, T) = \sum_{\mathbf{w} \in W} \left(\sum_{f \in \text{Hom}(FR(D, \mathbf{w}), T)} z^{srp_{\text{Im}(f) \subset T}^{m,n}(s, t)} \right)$$

and


$$rpp_{m,n}(L, T) = \sum_{\mathbf{w} \in W} \left(\sum_{f \in \text{Hom}(FR(D, \mathbf{w}), T)} \prod_{i=1}^c q_i^{w_i} z^{srp_{\text{Im}(f) \subset T}^{m,n}(s, t)} \right).$$

Specializing $s = t = 0$ (or, indeed, $z = 1$) in the subrack polynomial rpp yields the rack counting polynomial. Since every finite quandle T is a rack, the fact that subquandle polynomial invariants are stronger than unenhanced quandle counting invariants means *a fortiori* that subrack polynomial invariants are stronger than unenhanced rack counting invariants. The next example shows how subrack enhancement gives more information about a knot or link than the unadorned rack counting invariant.

Example 7 The trefoil knot 3_1 has simple rack counting invariant value 20 with respect to the rack with rack matrix below.



odd writhe



even writhe

$$M_T = \begin{bmatrix} 1 & 3 & 2 & 1 & 1 \\ 3 & 2 & 1 & 2 & 2 \\ 2 & 1 & 3 & 3 & 3 \\ 4 & 4 & 4 & 5 & 5 \\ 5 & 5 & 5 & 4 & 4 \end{bmatrix}$$

Here $N(T) = 2$, so we need only consider two diagrams of 3_1 , one with even writhe and one with odd writhe. The rack counting polynomial here is $11 + 9q$, which says that 11 colorings are contributed from the even-writhe diagram and 9 are contributed by the odd-writhe diagram. The subrack polynomial invariant is $rpp(3_1, T) = 2z^{2s^3t^3} + 3z^{s^3t^3} + 6z^{3s^3t^3} + 3qz^{s^3t^3} + 6qz^{3s^3t^3}$, which further filters the contributions – of the nine colorings of the odd-writhe diagram, six use colors in the subrack with subrack polynomial $3s^3t^3$ (in this case, the subquandle $\{1, 2, 3\}$) while three have colors in subracks with subrack polynomial s^3t^3 (here, the singleton subquandles $\{1\}, \{2\}, \{3\}$). Similarly, the even-writhe diagram has colorings corresponding to the odd-writhe colorings as expected, but additionally has two colorings by the subrack $\{4, 5\}$.

Indeed, the example suggests the following

Proposition 9 *If $S \subset T$ is a quandle and K a knot, then the contributions to $rpp(K, T)$ from S are equal for all powers of q . That is, rpp includes the term*

$$\left(\sum_{f \in \text{Hom}(Q(K), S)} z^{sp_{\text{Im}(f) \subset T}} \right) (1 + q + \dots + q^{N(T)-1}).$$

Proof. Quandle colorings do not depend on framing, so we get the same contribution, namely

$$\sum_{f \in \text{Hom}(Q(K), S)} z^{SP_{\text{Im}(f)} \subset T},$$

from each framing. □

6 Questions

We have shown that some classes of racks are classified by their generalized rack polynomials (the constant action racks) while others are not (quandles with polynomial ns^at^a). What conditions are sufficient for a type of rack to be determined by its generalized rack polynomials?

Since ns^at^a quandles are not determined by their generalized quandle polynomials, what extra information is necessary to determine these quandles up to isomorphism? How can such extra information be incorporated into the enhanced rack counting invariants?

For every rack R , the quotient rack under operator equivalence ($x \sim y \iff z \triangleright x = z \triangleright y \forall z$) is a quandle. What is the relationship between the subrack polynomial invariant with respect to R and the subquandle polynomial invariant with respect to $Q = R / \sim$?

References

- [1] N. Andruskiewitsch and M. Graña. From racks to pointed Hopf algebras. *Adv. Math.* **178** (2003) 177-243.
- [2] E. Brieskorn, Automorphic sets and braids and singularities. *Contemp. Math.* **78** (1988) 45-115.
- [3] R. Fenn and C. Rourke. Racks and links in codimension two. *J. Knot Theory Ramifications* **1** (1992) 343-406.
- [4] D. Joyce. A classifying invariant of knots, the knot quandle. *J. Pure Appl. Algebra* **23** (1982) 37-65.
- [5] S. V. Matveev. Distributive groupoids in knot theory. *Math. USSR, Sb.* **47** (1984) 73-83.
- [6] S. Nelson. A polynomial invariant of finite quandles. *J. Alg. Appl.* **2** (2008) 263-273.
- [7] S. Nelson. Generalized quandle polynomials. arXiv:0801.2979, to appear in *Can. Bull. Math.*
- [8] S. Nelson. Link invariants from finite racks. arXiv:0808.0029
- [9] S. Nelson and R. Wieghard. Link invariants from finite Coxeter racks. arXiv:0808.1584
- [10] H. Ryder. The structure of racks. PhD dissertation, U. Warwick (1993). Available at www.esotericka.org/quandles
- [11] M. Takasaki. Abstraction of symmetric transformation *Tohoku Math J.* **49** (1943) 145-207.

DEPARTMENT OF MATHEMATICS, POMONA COLLEGE, 610 N. COLLEGE AVE, CLAREMONT, CA 91711

Email address: tnc02005@mymail.pomona.edu

DEPARTMENT OF MATHEMATICS, CLAREMONT MCKENNA COLLEGE, 850 COLUMBIA AVE., CLAREMONT, CA 91711

Email address: knots@esotericka.org